# (19) World Intellectual Property Organization International Bureau



## 

## (43) International Publication Date 25 October 2001 (25.10.2001)

#### **PCT**

# (10) International Publication Number WO 01/78739 A1

- (51) International Patent Classification7: A61K 31/575, 31/46, A61P 11/06 // (A61K 31/575, 31:46)
- (21) International Application Number: PCT/GB01/01631
- (22) International Filing Date: 11 April 2001 (11.04.2001)
- (25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data: 0009606.5

18 April 2000 (18.04.2000) G

- (71) Applicant (for all designated States except US): GLAXO GROUP LIMITED [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): GAVIN, Brian, Charles [IE/IE]; GlaxoSmithKline, PO Box 700, Grange Road, Rathfarnfam, 16 Dublin (IE).
- (74) Agent: LEAROYD, Stephanie, Anne; GlaxoSmithKline, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



/78739 A

(54) Title: MEDICAL COMBINATIONS COMPRISING TIOTROPIUM AND FLUTICASONE PROPRIONATE

(57) Abstract: The present invention is concerned with pharmaceutical formulations comprising a combination of tiotropium and fluticasone propionate and the use of such formulations in medicine, particularly in the prophylaxis and treatment of respiratory diseases.

MEDICAL COMBINATIONS COMPRISING TIOTROPIUM AND FLUTICASONE PROPRIONATE

The present invention is concerned with combinations of tiotropium and fluticasone propionate, particularly compositions containing a combination of tiotropium and fluticasone propionate and the use of such compositions in medicine, particularly in the prophylaxis and treatment of respiratory diseases.

5

10

15

20

25

30

35

Tiotropium i.e.  $(1\alpha,2\beta,4\beta,5\alpha,7\beta)$ -7-[(hydroxydi-2-thienylacetyl)oxy]-9,9-dimethyl-3-oxa-9-azoniatricyclo[3.3.2.0]nonane and particularly its bromide salt is a well-known anti-cholinergic agent, described in EP418,716 for the treatment of bronchial asthma and related disorders.

Fluticasone propionate is an anti-inflammatory corticosteroid, described in GB 2088877, and is systematically named S-fluoromethyl-6α,9α-difluoro-11β-hydroxy-16α-methyl-17α-propionyloxy-3-oxoandrosta-1,4-diene-17β-carbothioate. Fluticasone propionate is now used clinically for the treatment of bronchial asthma and related disorders.

Although tiotropium bromide and fluticasone propionate may be effective therapies, there exists a clinical need for asthma therapies having potent and selective action and having an advantageous profile of action.

Therefore, according to the present invention there is provided a combination of tiotropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and fluticasone propionate or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof.

It will be appreciated that the compounds of the combination may be administered simultaneously, either in the same or different pharmaceutical formulations or sequentially. If there is sequential administration, the delay in administering the second compound should not be such as to lose the beneficial therapeutic effect of the combination.

According to a further aspect of the present invention, there is provided a pharmaceutical formulation comprising tiotropium or a pharmaceutically

2

PCT/GB01/01631

acceptable salt, solvate, or physiologically functional derivative thereof and fluticasone propionate or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients. According to a preferred aspect of the present invention, there is provided a pharmaceutical formulation comprising tiotropium bromide and fluticasone propionate, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients. In the most preferred aspect, the above pharmaceutical formulations are suitable for administration by

It is to be understood that the present invention covers all combinations of particular and preferred aspects of the invention described herein.

By the term "physiologically functional derivative" is meant a chemical derivative of tiotropium or fluticasone propionate having the same physiological function as the free compound, for example, by being convertible in the body thereto.

According to the present invention, examples of physiologically functional derivatives include esters.

20

25

30

35

5

10

WO 01/78739

inhalation.

Suitable salts according to the invention include those formed with both organic and inorganic acids. Pharmaceutically acceptable acid addition salts include but are not limited to those formed from hydrochloric, hydrobromic, sulphuric, citric, tartaric, phosphoric, lactic, pyruvic, acetic, trifluoroacetic, succinic, oxalic, fumaric, maleic, oxaloacetic, methanesulphonic, ethanesulphonic, p-toluenesulphonic, benzenesulphonic, isethionic, and naphthalenecarboxylic, such as 1-hydroxy-2-naphthalenecarboxylic acids.

Pharmaceutically acceptable esters of tiotropium or fluticasone propionate may have a hydroxyl group converted to a C<sub>1-6</sub>alkyl, aryl, aryl C<sub>1-6</sub> alkyl, or amino acid ester.

As mentioned above, both tiotropium and fluticasone propionate and their pharmaceutically acceptable salts, solvates, and physiologically functional derivatives have been described for use in the treatment of respiratory diseases.

3

Therefore, formulations of tiotropium and fluticasone propionate and their pharmaceutically acceptable salts, solvates, and physiologically functional derivatives have use in the prophylaxis and treatment of clinical conditions for which anticholinergic agent and/or an antiinflammatory corticosteroid is indicated. Such conditions include diseases associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary diseases (COPD) (e.g. chronic and wheezy bronchitis, emphysema), respiratory tract infection and upper respiratory tract disease.

5

10

15

20

25

30

35

Accordingly, the present invention provides a method for the prophylaxis or treatment of a clinical condition in a mammal, such as a human, for which an anticholinergic agent and/or antiinflammatory corticosteroid is indicated, which comprises administration of a therapeutically effective amount of a combination of tiotropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and fluticasone propionate or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof. The present invention further provides a method for the prophylaxis or treatment of a clinical condition in a mammal, such as a human, for which an anticholinergic agent and/or antiinflammatory corticosteroid is indicated, which comprises administration of a therapeutically effective amount of a pharmaceutical formulation comprising tiotropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and fluticasone propionate or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or excipient. In a preferred aspect, there is provided such a method which comprises administration of a therapeutically effective amount of a pharmaceutical formulation comprising tiotropium bromide and fluticasone propionate, and a pharmaceutically acceptable carrier or excipient. In particular, the present invention provides such methods for the prophylaxis or treatment of a disease associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary disease (COPD), respiratory tract infection or upper respiratory tract disease.

In the alternative, there is provided a combination of tiotropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and fluticasone propionate or a pharmaceutically acceptable salt,

WO 01/78739

4

PCT/GB01/01631

solvate, or physiologically functional derivative thereof, for use in therapy, particularly for use in the prophylaxis or treatment of a clinical condition for which an anticholinergic agent and/or antiinflammatory corticosteroid is indicated. In particular, there is provided a pharmaceutical formulation comprising tiotropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof (suitably, tiotropium bromide) and fluticasone propionate or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or excipient for use in therapy, particularly for use in the prophylaxis or treatment of a clinical condition for which an anticholinergic agent and/or antiinflammatory corticosteroid is indicated. In a preferred aspect, the invention is concerned with the prophylaxis or treatment of a disease associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary disease (COPD), respiratory tract infection or upper respiratory tract disease.

15

20

10

5

The amount of tiotropium and fluticasone propionate, or a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof which is required to achieve a therapeutic effect will, of course, vary with the particular compound, the route of administration, the subject under treatment, and the particular disorder or disease being treated. As a monotherapy, tiotropium bromide is generally administered to adult humans by aerosol inhalation at a dose of 10mcg to 200mcg twice daily. As a monotherapy, fluticasone propionate is administered to adult humans by aerosol inhalation at a dose of from 100mcg to 1000mcg twice daily, preferably 200mcg to 500mcg.

25

While it is possible for the active ingredients of the combination to be administered as the raw chemical, it is preferable to present them as a pharmaceutical formulation. When the individual compounds of the combination are administered separately, they are generally each presented as a pharmaceutical formulation as described previously in the art.

30

35

Pharmaceutical formulations are often prescribed to the patient in "patient packs" containing the whole course of treatment in a single package. Patient packs have an advantage over traditional prescriptions, where a pharmacist divides a patient's supply of a pharmaceutical from a bulk supply, in that the

5

patient always has access to the package insert contained in the patient pack, normally missing in traditional prescriptions. The inclusion of a package insert has been shown to improve patient compliance with the physician's instructions and, therefore, lead generally to more successful treatment. It will be understood that the administration of the combination of the invention by means of a single patient pack, or patient packs of each component compound, and containing a package insert instructing the patient to the correct use of the invention is a desirable additional feature of the invention.

Hereinafter, the term "active ingredients" means tiotropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, preferably tiotropium bromide, and fluticasone propionate, or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof.

Suitably, the pharmaceutical formulations which are suitable for inhalation according to the invention comprise the active ingredients in amounts such that each actuation provides therapeutically effective dose, for example, a dose of tiotropium of 10mcg to 200mcg, preferably 20mcg to 100mcg and a dose of fluticasone propionate of 50mcg to 1.0mg, preferably 100mcg to 500mcg.

20

25

15

5

The pharmaceutical formulations according to the invention may further include other therapeutic agents for example anti-inflammatory agents such as other corticosteroids (e.g. beclomethasone dipropionate, mometasone furoate, triamcinolone acetonide or budesonide) or NSAIDs (e.g. sodium cromoglycate, nedocromil sodium, PDE-4 inhibitors, leukotriene antagonists, iNOS inhibitors, tryptase and elastase inhibitors, beta-2 integrin antagonists and adenosine 2a agonists)) or,  $\beta_2$ -adrenoreceptor agonists (such as salbutamol, formoterol, salmeterol, fenoterol or terbutaline and salts thereof), or other anticholinergic agents (such as ipratropium).

30

35

The formulations include those suitable for oral, parenteral (including subcutaneous, intradermal, intramuscular, intravenous and intraarticular), intranasal, inhalation (including fine particle dusts or mists which may be generated by means of various types of metered dose pressurised aerosols, nebulisers or insufflators), rectal and topical (including dermal, buccal, sublingual

6

and intraocular) administration although the most suitable route may depend upon for example the condition and disorder of the recipient. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active ingredients into association with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

10

15

20

5

Formulations for inhalation include powder compositions which will preferably contain lactose, and spray compositions which may be formulated, for example, as aqueous solutions or suspensions or as aerosols delivered from pressurised packs, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, 1,1,1,2,3,3,3-heptafluoropropane, 1,1,1,2-tetrafluoroethane, carbon dioxide or other suitable gas. Suitable aerosol formulations include those described in EP 0372777 and WO93/11743. For suspension aerosols, the active ingredients should be micronised so as to permit inhalation of substantially all of the active ingredients into the lungs upon administration of the aerosol formulation, thus the active ingredients will have a particle size of less than 100 microns, desirably less than 20 microns, and preferably in the range 1 to 10 microns, for example, 1 to 5

25

microns.

Intranasal sprays may be formulated with aqueous or non-aqueous vehicles with the addition of agents such as thickening agents, buffer salts or acid or alkali to adjust the pH, isotonicity adjusting agents or anti-oxidants.

30 .

Capsules and cartridges or for example gelatin, or blisters of for example laminated aluminium foil, for use in an inhaler or insuflator may be formulated containing a powder mix of the active ingredients and a suitable powder base such as lactose or starch. In this aspect, the active ingredients are suitably micronised so as to permit inhalation of substantially all of the active ingredients into the lungs upon administration of the dry powder formulation, thus the active

7

ingredients will have a particle size of less than 100 microns, desirably less than 20 microns, and preferably in the range 1 to 10 microns.

Solutions for inhalation by nebulation may be formulated with an aqueous vehicle with the addition of agents such as acid or alkali, buffer salts, isotonicity adjusting agents or antimicrobials. They may be sterilised by filtration or heating in an autoclave, or presented as a non-sterile product.

Preferred unit dosage formulations are those containing a pharmaceutically
effective dose, as hereinbefore recited, or an appropriate fraction thereof, of the
active ingredient. Thus, in the case of formulations designed for delivery by
metered dose pressurised aerosols, one actuation of the aerosol may deliver
half of the therapeutically effective amount such that two actuations are
necessary to deliver the therapeutically effective dose.

15

20

5

It should be understood that in addition to the ingredients particularly mentioned above, the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question. Furthermore, the claimed formulations include bioequivalents as defined by the US Food and Drugs Agency.

For a better understanding of the invention, the following Examples are given by way of illustration.

#### 25 EXAMPLES

#### A: Metered Dose Inhalers

#### Example 1

	Per actuation	
tiotropium bromide	100 microgram	
fluticasone propionate	200 microgram	
1,1,1,2-Tetrafluoroethane	nane to 75.0mg	

8

The micronised active ingredients are weighed into an aluminium can, 1,1,1,2-tetrafluoroethane is then added from a vacuum flask and a metering valve is crimped into place.

•5

Similar methods may be used for the formulation of Examples 2 to 4:

## Example 2

	Per actuation	
tiotropium bromide	200 microgram	
fluticasone propionate	100 microgram	
1,1,1,2-Tetrafluoroethane	to 75.0mg	

10

## Example 3

	Per actuation	
tiotropium bromide	18 microgram	
fluticasone propionate	100 microgram	
1,1,1,2-Tetrafluoroethane	to 75.0mg	

## Example 4

15

	Per actuation	
tiotropium bromide	mide 9 microgram	
fluticasone propionate	100 microgram	
1,1,1,2-Tetrafluoroethane	to 75.0mg	

9

## **B**: Dry Powder Inhalers

## Example 5

	Per cartridge or blister
tiotropium bromide	100 microgram
fluticasone propionate	200 microgram
Lactose Ph. Eur.	to 12.5mg
	or to 25.0mg

5

The active ingredients are micronised and bulk blended with the lactose in the proportions given above. The blend is filled into hard gelatin capsules or cartridges or in specifically constructed double foil blister packs to be administered by an inhaler such as a Rotahaler, Diskhaler, or Diskus inhaler (each of these being a Trademark of Glaxo Group Limited).

Similar methods may be used for the formulations of Examples 6 to 8:

#### Example 6

15

10

	Per cartridge or blister
tiotropium bromide	200 microgram
fluticasone propionate	100 microgram
Lactose Ph. Eur.	to 12.5mg
	or to 25.0mg

## Example 7

	Per cartridge or blister	
tiotropium bromide	18 microgram	
fluticasone propionate	100 microgram	
Lactose Ph. Eur.	to 12.5mg	
	or to 25.0mg	

10

## Example 8

	Per cartridge or blister	
tiotropium bromide	9 microgram	
fluticasone propionate	100 microgram	
Lactose Ph. Eur.	to 12.5mg	
	or to 25.0mg	

#### **Claims**

5

- A pharmaceutical formulation comprising tiotropium or a
   pharmaceutically acceptable salt, solvate, or physiologically functional
   derivative thereof and fluticasone propionate or a pharmaceutically
   acceptable salt, solvate, or physiologically functional derivative thereof,
   and a pharmaceutically acceptable carrier or excipient, and optionally
   one or more other therapeutic ingredients.
- 10 2. A pharmaceutical formulation comprising tiotropium bromide and fluticasone propionate, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients.
- 3. A pharmaceutical formulation according to claim 1 or 2 which is suitable for administration by inhalation.
  - 4. A pharmaceutical formulation according to any of claims 1 to 3 wherein the pharmaceutically acceptable carrier or excipient is lactose.
- 20 5. A pharmaceutical formulation according to any of claims 1 to 3 wherein the pharmaceutically acceptable carrier or excipient comprises 1,1,1,2-tetrafluoroathane and/or 1,1,1,2,3,3,3-heptafluoropropane.
- 6. A method for the prophylaxis or treatment of a clinical condition in a mammal, such as a human, for which an anticholinergic agent and/or antiinflammatory corticosteroid is indicated, which comprises administration of a therapeutically effective amount of a pharmaceutical formulation according to any one of claims 1 to 5.
- 30 7. A method according to claim 6 wherein the clinical condition is a disease associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary disease (COPD), respiratory tract infection or upper respiratory tract disease.

	INTERNATIONAL SEARCH REPORT	In 181 Application No PCT/GB 01/01631
A. CLASSI IPC 7	FICATION OF SUBJECT MATTER A61K31/575 A61K31/46 A61P11/06 //(A61K	31/575,31:46)
According to	International Patent Classification (IPC) or to both national classification and IPC	
	SEARCHED	
IPC 7	cumentation searched (classification system followed by classification symbols)  A61K A61P	
	ion searched other than minimum documentation to the extent that such documents are incl	
}	ala base consulted during the International search (name of data base and, where practica ternal, WPI Data, MEDLINE, BIOSIS, CHEM ABS Data	•
	ENTS CONSIDERED TO BE RELEVANT	
Category °	Citation of document, with Indication, where appropriate, of the relevant passages	Relevant to claim No.
Υ	EP 0 418 716 A (BOEHRINGER INGELHEIM KG;BOEHRINGER INGELHEIM INT (DE)) 27 March 1991 (1991-03-27) cited in the application claims	1-7
Υ	QURESHI F ET AL: "Effect of nebulized ipratropium on the hospitalization rates of children with asthma."  NEW ENGLAND JOURNAL OF MEDICINE, (1998 OCT 8) 339 (15) 1030-5.,  XP001007631  abstract /	1-7
X Furti	ner documents are listed in the continuation of box C. X Patent family	members are listed in annex.
° Special ca	tegories of cited documents : "T" later document put	olished after the international filing date
	ent defining the general state of the art which is not cited to understate ered to be of particular relevance	olished after the international filing date d not in conflict with the application but nd the principle or theory underlying the

<ul> <li>Special categories of cited documents:</li> <li>"A" document defining the general state of the art which is not considered to be of particular relevance</li> <li>"E" earlier document but published on or after the international filling date</li> <li>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</li> <li>"O" document referring to an oral disclosure, use, exhibition or other means</li> <li>"P" document published prior to the international filing date but later than the priority date claimed</li> </ul>	<ul> <li>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</li> <li>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</li> <li>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</li> <li>"&amp;" document member of the same patent family</li> </ul>
Date of the actual completion of the international search  8 August 2001	Date of mailing of the international search report $04/09/2001$
Name and mailing address of the ISA  European Patent Office, P.B. 5818 Patentiaan 2  NL – 2280 HV Rijswijk  Tel. (+31–70) 340–2040, Tx. 31 651 epo nl, Fax: (+31–70) 340–3016	Authorized officer  Herrera, S

#### INTERNATIONAL SEARCH REPORT

In nal Application No
PCT/GB 01/01631

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT			
Category •	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	
Y	BACULARD A: "Bronchodual in the long-term treatment of children with asthma." ARCHIVES DE PEDIATRIE, vol. 2, no. SUPPL. 2, 1995, pages 149S-153S, XP000914115 ISSN: 0929-693X abstract	1-7	
Y	GB 2 088 877 A (GLAXO GROUP LTD) 16 June 1982 (1982-06-16) cited in the application claims 1,20-24	1~7	
Y	BOWLER S: "LONG ACTING BETA AGONISTS" AUSTRALIAN FAMILY PHYSICIAN, XX, XX, vol. 27, no. 12, December 1998 (1998-12), pages 1115,1117-1118, XP000973076 the whole document	1-7	
Y	O'CONNOR B J: "COMBINATION THERAPY" PULMONARY PHARMACOLOGY AND THERAPEUTICS, ACADEMIC PRESS, NEW YORK, NY, US, vol. 11, no. 5/6, 1998, pages 397-399, XP000911059 ISSN: 1094-5539 the whole document	17	

## INTERNATIONAL SEARCH REPORT

iformation on patent family members

II PCT/GB 01/01631

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0418716 A	27-03-1991	DE 3931041 A AT 103914 T AU 642913 B AU 6431890 A BG 61295 B CA 2066248 A,C CZ 9004523 A DD 297647 A DE 59005250 D DK 418716 T WO 9104252 A ES 2052125 T HR 940723 A HU 60740 A HU 208823 B HU 210612 B IE 903342 A IL 95691 A JP 7030074 B JP 5502438 T KR 168432 B MX 9203150 A NO 301478 B NZ 235306 A PL 168468 B PT 95312 A,B	28-03-1991 15-04-1994 04-11-1993 18-04-1997 17-03-1991 11-11-1998 16-01-1992 11-05-1994 02-05-1994 04-04-1991 01-07-1994 30-06-1997 28-10-1992 28-01-1994 29-05-1995 10-04-1991 23-07-1996 05-04-1995 28-04-1993 15-01-1999 01-07-1992 03-11-1997 24-06-1997 29-02-1996 22-05-1991
GB 2088877 A	16-06-1982	SI 9011744 A,B SK 452390 A RU 2073677 C US 5610163 A ZA 9007338 A  AT 395428 B AT 17084 A AT 401521 B AT 34491 A AT 395427 B AT 395429 B AT 203186 A AU 544517 B AU 6729881 A BE 887518 A BG 60700 B CA 1201114 A CA 1205464 C CH 644615 A CH 651307 A CZ 9104034 A CY 1291 A DE 3105307 A DE 3153379 C DK 62381 A,B, ES 499394 D ES 8207194 A ES 509539 D ES 8305379 A	31-10-1997 04-11-1998 20-02-1997 11-03-1997 26-08-1992 25-08-1992 25-09-1996 15-02-1996 28-12-1992 25-09-1996 28-12-1992 28-12-1992 28-12-1992 28-12-1992 28-12-1992 28-12-1992 28-12-1995 20-08-1981 13-08-1981 29-12-1995 25-02-1986 03-06-1986 15-08-1984 13-09-1985 16-03-1994 18-10-1985 10-12-1981 19-11-1992 16-08-1981 01-09-1982 01-12-1982 01-04-1983 01-07-1983

## INTERNATIONAL SEARCH REPORT

iformation on patent family members

Ini 1al Application No
PCT/GB 01/01631

	<del>,</del>			<del>,                                      </del>
Patent document cited in search report	Publication date		Patent family member(s)	Publication date
GB 2088877 A		ES	8402317 A	16-04-1984
		ES	524985 D	01-01-1985
		ES	8502447 A	01-04-1985
		ES	532055 D	16-10-1985
		ES	8600936 A	16-02-1986
		FI	810444 A,B,	16-08-1981
		FR	2477156 A	04-09-1981
		FR	2485542 A	31-12-1981
	•	GB	2137206 A,B	03-10-1984
		HK	58385 A	16-08-1985
		ΙE	51394 B	24-12-1986
		ΙE	51395 B	24-12-1986
		IT	1170717 B	03-06-1987
		JP	1488353 C	23-03-1989
		JP	56138200 A	28-10-1981
		JP	63037120 B	22-07-1988
		KE	3526 A	07-06-1985
		KR	8500969 B	02-07-1985
		MX	9202717 A	30-06-1992
		MY	75785 A	31-12-1985
		NL	84649 C	
		NL	8100707 A,B,	16-09-1981
		NZ	196260 A	30-11-1983
		PH	24267 A	29-05-1990

Form PCT/ISA/210 (patent family annex) (July 1992)